

# Conformal Mapping Using the Schwarz-Christoffel Transform 2007-2008

Evan Warner

November 14, 2007

## Abstract

The Schwarz-Christoffel transform is a conformal mapping from the upper half of the complex plane to a polygonal domain. It allows many physical problems posed on two-dimensional, polygonal regions, such as heat flow, fluid flow, and electrostatics, to be solved numerically. This type of problem cannot generally be solved in closed form; the Schwarz-Christoffel transform provides an exceptionally accurate method of solution. This project will produce a working software unit that efficiently and accurately calculates Schwarz-Christoffel transforms and inverses. The program will incorporate graphical, easy-to-use interfaces and contain resources to aid in solving physical problems. In addition, research into mathematical extensions to the Schwarz-Christoffel transform, such as the inclusion of simple curves, will be conducted.

## Introduction

Many physical problems are expressed as differential or boundary value problems over a surface. Often, these surfaces are or can be approximated by two-dimensional polygons. In this specific case, one method of determining accurate solutions is by taking the polygonal domain to exist in the complex plane and determining a conformal map, which preserves the structure of Laplace's equation, that restates the problem in a simpler domain, most often the upper half-plane. The new problem, now easy to solve analytically or in closed form, is then mapped back to the original domain. For such polygonal domains, a method of determining the specific transform needed is provided by the following formula, known as the Schwarz-Christoffel transform:

$$f(z) = A \int_0^z \prod_{j=1}^n (\zeta - x_j)^{-\theta_j/\pi} d\zeta + B. \quad (1)$$

In this formula,  $\zeta$  is an independent complex variable in the upper half-plane, the  $\theta_j$  are the exterior angles of the polygon, the  $x_j$  are 'prevertices' of the mapping (given along the real axis),  $n$  is the number of vertices of the polygon, and  $A$  and  $B$  are complex constants that specify the location of the image polygon in the complex plane. The  $\theta_j$  must satisfy

$$\sum_{j=1}^n \theta_j = 2\pi, \quad (2)$$

which ensures the completeness of the image polygon [2]. Unfortunately, the Schwarz-Christoffel formula is not easy to evaluate, and requires both effective integration algorithms and efficient, convergent nonlinear equation solvers. Implementation of such numerical routines is not a trivial problem, and is the subject of this paper.

## Background

The Schwarz-Christoffel transform was first discovered independently in the late 1860s by Elwin Christoffel and Hermann Schwarz. Schwarz used some of the ideas of the transform to provide a more rigorous proof of the Riemann Mapping Theorem, which he had previously shown to be incomplete, but the majority of this work was on a purely theoretical level [5]. The usefulness of the transform was mitigated by the formula's unwieldiness, as the mappings for all but the simplest domains could not be calculated in closed form. Numerical estimates, especially for nonsymmetric polygons with four or more vertices, could not be effectively calculated by hand. Application to physical problems, therefore, was limited at best until the advent of the computer. A computer algorithm to compute the Schwarz-Christoffel transform was first written in the 1960s, and others have been written and modified since then [3].

The first problem in calculating the Schwarz-Christoffel mapping is the evaluation of the integral given by Eq. (1). The integrand contains singularities at each of the endpoints of the image polygon, which tend to render ordinary numerical integration routines either useless or hopelessly slow. In addition, the presence of negative powers in  $f$  means that domains of applicability for each of the subfunctions  $(\zeta - x_j)^{-\theta_j/\pi}$  must be chosen so that the entire domain in and immediately around the image polygon is meromorphic. Although several quadrature routines have been used for this problem, the method of choice today is Gauss-Jacobi quadrature, which uses a specially-tailored weighting function to choose points of evaluation and weights for the points that maximize efficiency. In practice, the Schwarz-Christoffel formula is altered so that the prevertex  $x_n$  is chosen to be both  $-\infty$  and  $+\infty$  (the values are equivalent for a conformal map, which acts on the Riemann sphere). This can always be done due to the extra degrees of freedom contained in Eq. (1). The integrals that must be evaluated in practice in the course of the Schwarz-Christoffel transform are of the form

$$\int_{x_{i-1}}^{x_i} \prod_{j=1}^{n-1} (\zeta - x_j)^{-\theta_j/\pi} d\zeta. \quad (3)$$

These integrals can always be written as required for Gauss-Jacobi quadrature; that is, in the form

$$\int_a^b (z-a)^\alpha (z-b)^\beta \psi(z) dz, \quad (4)$$

where  $\alpha$  and  $\beta$  are real numbers greater than  $-1$ .

The points and weights of a Gauss-Jacobi quadrature are calculated here using a routine from Numerical Recipes [4] which efficiently estimates and solves for the roots of the Jacobi polynomials, which form the sample points just as the roots of the Chebyshev polynomials form the sample points for standard Gaussian quadrature. These points, however, are uniformly calculated in the range  $[-1, 1]$ , and the integrals must be adjusted slightly to conform to this range. During the calculation of the prevertices, discussed below, the  $z$  in Eq. (4) will be restricted to the real axis; however, in direct calculations once the prevertices have been found, the  $z$  will generally be fully complex, which must be dealt with by the program.

The second problem is the Schwarz-Christoffel parameter problem, where the  $x_j$  in Eq. (1) are calculated. As described in [2], a series of nonlinear, constrained equations can be formed from the requirement that the image polygon and the desired polygon be similar (the constants  $A$  and  $B$  in Eq. (1) then ensure congruency). Written out, there are  $n-3$  linear equations in  $n-3$  unknowns, once the extra degrees of freedom have been taken care of by arbitrarily giving three of the  $x_j$  precise values. Here, as in the literature, we take  $x_1 = -1$  and  $x_2 = 0$  in addition to the already-defined  $x_n = \pm\infty$ . The equations to be solved are then

$$\frac{|\int_{x_{i-1}}^{x_i} \prod_{j=1}^n (\zeta - x_j)^{-\theta_j/\pi} d\zeta|}{|\int_{x_1}^{x_2} \prod_{j=1}^n (\zeta - x_j)^{-\theta_j/\pi} d\zeta|} = \frac{|w_j - w_{j-1}|}{|w_2 - w_1|} = 0, \quad (5)$$

where  $i = 3, 4, \dots, n-1$ . However, there is an additional complication, as the order of the prevertices on the real axis matters. The extra constraint can be expressed as

$$0 < x_3 < x_4 < \dots < x_{n-1} < \infty. \quad (6)$$

The unconstrained problem is relatively easy to solve; however, the constraint prevents a naive application of a Newton's Method variant to this problem. To get around this, Trefethen in [3] suggests a simple change of variables that ensures the inequalities of Eq. (6). Take a new series of variables,  $\chi_j$ , and let

$$\chi_j = \ln x_j - x_{j-1}. \quad (7)$$

The resulting  $\chi_j$  will automatically obey Eq. (6), and the original  $x_j$  are found by the simple inverse formula

$$x_j = x_{j-1} + e^{\chi_j}. \quad (8)$$

This new set of equations in the  $\chi_j$  is readily solved by a variant of Newton's Method that does not require the calculation of the Jacobian matrix (which would be hopelessly complex), but rather uses progressive estimates.

## Development

So far, the software has been written entirely in Java, although certain routines may be later written in C to increase speed if there is a bottleneck at any point in the process. The entire development of the program is designed to be achieved in stages by attacking the subproblems individually. The following is a list of classes, with short descriptions, written up to this point:

- **class Complex** - this class stores and performs arithmetic on complex numbers, which are not directly supported by Java. Several of the methods, including the multiplication and division algorithms, are designed to run as quickly as possible while avoiding intermediate overflow and floating-point error propagation. The multiplication method, for instance, requires only three real multiplications rather than four.
- **class ComplexFunction** - this class stores and evaluates functions of one complex variable. Its constructor accepts a String argument and stores the function, if syntactically acceptable, as a String in postfix form. The function may then be evaluated at any point on the complex plane. (see *Appendix*)
- **class GaussJacobiWeights** - this class calculates and stores the sample points and weights for a given Gauss-Jacobi quadrature over the interval  $[-1, 1]$ . This routine uses Newton's Method to find the roots of the Jacobi polynomials, which are the sample points for the integral, and was taken and translated from [4].
- **class GaussQuad** - this class accepts as input  $\psi$ ,  $a$ ,  $b$ ,  $\alpha$ , and  $\beta$  from Eq. (4). For an arbitrary integral in that form, shifting and scaling the bounds produces the equivalent integral

$$c^{\alpha+\beta+1} \int_{-1}^1 (\zeta-1)^\alpha (\zeta+1)^\beta \psi(c\zeta+m) d\zeta, \quad (9)$$

where  $b = \frac{a+b}{2}$  and  $c = b-m = m-a$ . This integral is then evaluated using the sample points and weights given by the `GaussJacobiWeights` class and returned. For any `GaussQuad` object, varying numbers of sample points (and thus varying accuracy) are accepted by its `integrate()` method.

- **class SchwarzChristoffel** - this class runs the graphical user interface and calls `GaussQuad` when necessary. The graph itself has the ability to show axes and manually adjust window parameters.

Once the basic transform is operational, a new set of routines will be implemented to calculate continuous Schwarz-Christoffel problems. Immediately following from Eq. (3) above, we have

$$f'(z) = A \prod_{j=1}^{n-1} (\zeta - x_j)^{-\theta_j/\pi}. \quad (10)$$

To change this into a continuous problem, we can rewrite this as

$$f'(z) = A e^{\frac{1}{\pi} \sum_{j=1}^{n-1} -\theta_j \ln(\zeta - x_j)}. \quad (11)$$

Then, defining the natural logarithm function as single-valued in the upper half-plane, except where  $x_i = z$ ,  $f'$  becomes an analytic function in the required domain. To formulate the continuous-boundary problem, we simply replace the sum in Eq. (11) with an integral, and integrate the entire function to find  $f(z)$ :

$$f(z) = A \int_0^z e^{\frac{1}{\pi} \int_{-\infty}^{\infty} -\theta(x) \ln(\zeta - x_j) dx} d\zeta + B, \quad (12)$$

where  $\theta(x)$  represents the amount of turning per unit length on the real axis, such that

$$\int_{-\infty}^{\infty} \theta(x) dx = 2\pi. \quad (13)$$

The continuous problem therefore has an extra subproblem to solve, namely, the solution of the integral equation, Eq. (12), to find  $\theta(x)$  at every  $x$ . This will be dealt with in future iterations of the project.

## Expected Results

The purpose of this project was to calculate and display Schwarz-Christoffel transforms, which conformally map the upper half-plane to an arbitrary polygon, efficiently and accurately. In addition, additional research into the Schwarz-Christoffel transform itself, including its extension to curved target domains, was investigated. The evaluation of the Schwarz-Christoffel formula involves several parts, including the efficient calculation of a certain class of integrals as well as a solver of nonlinear systems of equations. Solving the continuous-parameter problem will require numerical solutions to a certain class of integral equations.

For a first-quarter version of the project, the software is moderately successful. The first problem, that of numerical integration, has been solved and refined, and a basic user interface has been designed. Once the equation solver has been implemented, the program itself will be fully functional on a basic level. The completed program will be useful on several levels: as a teaching aid, and as a tool for researchers solving certain equations on polygonal regions. Once the basic Schwarz-Christoffel problem is numerically solved, the program can form an easy basis for testing research in numerical analysis and mathematics that deals with improving or expanding the Schwarz-Christoffel transform.

## References

- [1] Howell, L. H. (1990). *Computation of conformal maps by modified Schwarz-Christoffel transformations*. Retrieved September 28, 2007, from <http://citeseer.ist.psu.edu/howell90computation.html>.
- [2] Saff, E. B., & Snider, A. D. (n.d.). *Fundamentals of complex analysis with applications to engineering, science, and mathematics*. Prentice-Hall Engineering/Science/Mathematics.
- [3] Trefethen, L. (1979). *Numerical computation of the Schwarz-Christoffel transformation*. Retrieved September 28, 2007, from <ftp://reports.stanford.edu/pub/cstr/reports/cs/tr/79/710/CS-TR-79-710.pdf>.
- [4] Press W., Teukolsky, S., Vetterling, W., & Flannery, B. (1992). *Numerical Recipes in C, Second Edition*. Cambridge University Press.
- [5] O'Connor, J.J., & Robertson, E.F. (2001). *Hermann Amandus Schwarz*. Retrieved November 3, 2007, from <http://www-history.mcs.st-andrews.ac.uk/Biographies/Schwarz.html>.